


LICE... LICE... LICE...

ALL THERE IS TO KNOW ABOUT HEAD LICE



Lice have been around for a long time
and they are not about to disappear any time soon.
They are not dangerous, although they can be annoying.




Did You Know

- Head lice are often found in children who go to school or to a child care centre, especially after summer vacation.
- They affect anybody, rich or poor.
- They don't care if your hair is dirty or clean, they still like it.
- They don't always cause itching.
- They aren't treated by washing your hair with regular shampoo: special louse treatment is necessary.
- They don't carry disease.

● ● ● Head lice

- They are minute greyish insects.
- They feed on human blood.
- They live near the hair root and can sometimes be found on eyebrows and in beards.
- They don't live on pets.
- They don't jump or fly.
- Females lay an average of 5 eggs (nits) per day.



••• Nits

- They cling to the hair shaft.
- They are difficult to remove.

LIVE NITS

- They are greyish-white eggs that look like puffy, shiny, and translucent dandruff flakes.
- They are usually found within 6 mm of the scalp.

DEAD NITS

- They are whiter and dried out.
- They are mainly found more than 6 mm away from the scalp.



 **A drake dandruff, dust, or dirt that can be easily removed is not a nit.**

••• How do you get head lice?

- Lice can spread **easily** through **head-to-head contact**, like a hug, for example.
- Head lice are **not easily spread through contact with personal belongings** such as hats, combs, hairbrushes, and pillowcases, or through contact with the back of sofas, or with sheets, carpets, eyeglasses, etc.



How do you find out if you have lice or nits?

By checking the head, preferably with a nit comb, a very fine-toothed comb designed for this purpose and available in pharmacies. Most often, there are less than 10 or 20 lice on an infested head, so it is important to perform a thorough examination.

WHEN?

- At least once a week, after shampooing, especially right after summer vacation.
- Every day, if there is a head lice infestation around.
- Whenever the scalp starts to get itchy.

HOW?


- Place the child's head under a good light: head lice flee from light and you can see them moving (a magnifying glass can help).
- Wet the hair, then comb it with the nit comb. Conditioner makes louse hunting easier. However, hair treatment products are less effective if you use a conditioner, silicone-based products or vinegar in the hair.
- Look for lice and nits, especially behind the ears and near the back of the neck.
- Separate the hair into tufts the width of the comb. In the case of very long hair, the tufts should be about 2 cm in width.
- Comb the hair from scalp to tip with a nit comb (fine tooth comb).
 - Examine the hair section by section, from one side to the other and from the forehead back to the nape of the neck.
 - Look for lice or nits on the comb each time you run it through the hair.



Wash your hands after each examination.

If you find lice or nits

- Consult a pharmacist as soon as possible so that he or she can prescribe the most appropriate head lice treatment.
- Apply head lice medication as soon as possible.
- Inform the school or child care centre and anyone who has been in close contact with the infested child, as soon as possible. As a rule, children should not have to stay home from school or daycare.
- **Check** the heads of **all other family members**.



Treat ONLY those who have lice or live nits. The treatment is more effective when all family members who are infested are treated **at the same time** (on the same day). It may be difficult to determine whether the nits are dead or alive. If in doubt, consult a healthcare professional before starting treatment.

How to prevent head lice

Remind children to:

- Avoid head-to-head contact.
- Keep hats, caps, and scarves in their coat sleeves.
- Not to share personal items, such as combs, hairbrushes, barrettes, and hats.
- Keep long hair tied back.

Check the child's head regularly.

There is no preventive treatment. Only early screening and treatment of those infested can prevent head lice from spreading.

Parents' cooperation is essential if head lice are to be prevented and controlled.

Treating

head lice



A person with live lice or nits (eggs) should be treated promptly with head-lice medication. Products for animals should not be used on humans.

• • • Choosing the product

Although head lice treatments are available over-the-counter, it is preferable to consult a pharmacist, who will be able to determine the most appropriate treatment.

Some treatments can be used by people who are pregnant or nursing persons. Ideally, before using head-lice medication, it is suggested that they consult their doctor or pharmacist.

Application of these products may cause itching or a mild burning sensation of the scalp.

If eyebrows are infested, apply petroleum jelly (Vaseline® or Lacri-Lube®) at least 2 to 4 times a day for 1 week. The dead nits can then be removed with the fingernails or tweezers, by sliding them along the eyebrow.



• • • Application principles

Depending on the treatment, two or three applications are needed on a specific treatment schedule, which the pharmacist will explain. Administer the recommended number of applications, even if you find no live lice or viable nits in the hair. For the best chance of success, use a nit comb to remove lice and nits during treatment.

If live lice are observed 17 days **after the first application of the product, or if there is any doubts about the treatment's effectiveness, a health professional (e.g., pharmacist)** should be consulted to evaluate the situation and advise on the appropriate course of action to get rid of the lice. Treatment failure could result from not following the treatment (e.g., not following the instructions or number of applications recommended) or because the lice are resistant to the product used. You could also have a new infestation.

How to apply:

Follow the manufacturer's instructions for the product selected.

When the treatment is finished according to the manufacturer's instructions, dry the hair with a fresh towel or a hair dryer. Put used towels in with dirty laundry.

Use a nit comb to remove lice and nits 2, 11, and 17 days after the first application. Using a nit comb on wet hair increases your chances of successful treatment

● ● ● **Removing nits**

- Apply a damp towel to the scalp for at least 30 minutes to help dislodge the nits.
- Place the head under a good light; a magnifying glass can help.
- Get a nit comb and work near a source of hot water.
- Untangle hair with an ordinary comb.
- Separate the hair into tufts the width of the comb. In the case of very long hair, the tufts should be about 2 cm in width.
- Comb through each tuft of hair with a nit comb.
- Angle the nit comb towards the scalp and insert it into the tuft as close to the scalp as possible.
- Make sure that the entire length of the teeth is in the tuft.
- Comb out along the complete length of the hair.
- Examine the hair section by section, from one side to the other and from the forehead back to the nape of the neck.
- Pay particular attention to behind the ears and the nape of the neck.
- Soak the nit comb in hot water after running it through each tuft of hair.

To remove a nit that the nit comb doesn't get, use fingernails or eyebrow tweezers to slide it along the length of the hair.

Both the ordinary and nit combs must be cleaned using one of the methods described on the next page.

• • • Washing personal belongings

The risk of infestation through contact with personal belongings is minimal, but if a notice from the school or child care centre is received, treat personal belongings (bedclothes, combs, hairbrushes, hats, caps, and other accessories) used on a regular basis by the infested person using one of the following methods:

Soak belongings in undiluted head-lice medication for 5 to 10 minutes.

or

Soak belongings in hot water (around 65°C or 150°F) for 5 to 10 minutes.

or

Put clothing in the dryer set at the warmest setting for 20 minutes.

or

Dry-clean clothing.

or

Store belongings in a sealed plastic bag for 10 days.



Spraying the house, furniture, or other household objects with pesticides is NOT recommended. Not only is this ineffective in controlling head lice, but it can also pose a serious health risk to people or animals exposed to a potentially toxic product. Carpets and sofas can be vacuumed.

This document may be found in the **Documentation** section of the Ministère de la Santé et des Services sociaux Web site at: Quebec.ca

Legal deposit
Bibliothèque et Archives nationales du Québec, 2023
ISBN 978-2-550-94820-9 (print version)
ISBN 978-2-550-94821-6 (PDF)

All rights reserved for all countries. Any reproduction whatsoever, translation or dissemination, in whole or in part, is prohibited unless authorized by Les Publications du Québec. However, reproduction in whole or in part for personal, non-commercial purposes is permitted solely in Québec, provided the source is mentioned.